

JCAHO CORE MEASURES

Acute MI	3Q09	JCAHO
Aspirin at arrival	98%	98%
Aspirin Prescribed at discharge	94%	98%
ACEI or ARB for LVSD	93%	94%
Adult smoking cessation advice / counseling	100%	99%
Beta Blocker prescribed at discharge	98%	98%
Mean time to PCI(minutes)	65	91
within 90(changed 3Q06) minutes of hospital arrival	93%	82%
Inpatient mortality	5%	5%
Heart Failure	3Q09	JCAHO
Discharge instructions	72%	82%
LVF assessment	100%	97%
ACEI or ARB for LVSD	94%	93%
Adult smoking cessation advice/counseling	100%	97%
Pneumonia	3Q09	JCAHO
Pneumococcal vaccination	91%	90%
Blood culture before first antibiotic	86%	93%
Adult smoking cessation advice / counseling	97%	95%
Antibiotic Timing (Mean)(minutes)	130	171
antibiotic received within 6 hours of hospital arrival	86%	93%
selection for CAP in immunocompetent - ICU patient	60%	60%
selection for CAP immunocompetent - Non ICU patient	96%	93%
Influenza vaccination	83%	83%
STROKE	3Q09	JCAHO
DVT Prophylaxis by end of hospital day two	100%	85%
Discharged on Antithrombotic therapy	92%	85%
Atrial Fibrillation received Anticoagulation Therapy	85%	85%
Arrive w/in 120min & IV t-PA initiated w/in 180min	25%	85%
Antithrombotic therapy by end of hospital day two	77%	85%
Discharged on Cholesterol Reducing drugs	100%	85%
Dysphagia Screening	97%	85%
Educational material provided	84%	85%
Smoking cessation advice/ counseling	100%	85%
Assessed for Rehabilitation Services	97%	85%

Core Measures

SCIP	3Q09	JCAHO
Prophylactic abx within 1 hr prior to surgical incision - Overall Rate	97%	92%
1hr prior to surgical incision - CABG	100%	93%
1hr prior to surgical incision - Other Cardiac Surgery	100%	93%
1hr prior to surgical incision - Hip Arthroplasty	100%	92%
1hr prior to surgical incision - Knee Arthroplasty	97%	95%
1hr prior to surgical incision - Colon Surgery	82%	87%
1hr prior to surgical incision - Hysterectomy	100%	93%
1hr prior to surgical incision - Vascular Surgery	100%	89%
Prophylactic abx selection surgical patients - Overall Rate	98%	96%
Abx selection surgical patients - CABG	100%	99%
Abx selection surgical patients - Other Cardiac Surgery	100%	99%
Abx selection surgical patients - Hip Arthroplasty	100%	99%
Abx selection surgical patients - Knee Arthroplasty	100%	99%
Abx selection surgical patients - Colon Surgery	91%	84%
Abx selection surgical patients - Hysterectomy	100%	96%
Abx selection surgical patients - Vascular Surgery	92%	96%
Prophylactic abx discontinued within 24hrs after surgery end time - Overall Rate	86%	89%
Discontinued within 48hrs after surgery - CABG	100%	93%
Discontinued within 48hrs after surgery - Other Cardiac Surgery	100%	92%
Discontinued within 24hrs after surgery - Hip Arthroplasty	82%	89%
Discontinued within 24hrs after surgery - Knee Arthroplasty	84%	90%
Discontinued within 24hrs after surgery - Colon Surgery	90%	80%
Discontinued within 24hrs after surgery - Hysterectomy	93%	92%
Discontinued within 24hrs after surgery - Vascular Surgery	56%	82%
Cardiac Surgery Patients with Controlled 6 a.m. Postoperative Serum Glucose	92%	89%
Surgery Patients with Appropriate Hair Removal	100%	97%
Colorectal Surgery Patients with Immediate Postoperative Normothermia	91%	96%
Surgery Pt on BB Therapy Received BB during Preoperative Period	100%	93%
Surgery Pt with Recommended Venous Thromboembolism Prophylaxis Ordered	80%	92%
Received Appropriate VTE Prophylaxis within 24 hrs Prior to thru 24hrs after Surgery	84%	89%